

Arshad, M.N.<sup>a</sup>, Khan, I.U.<sup>b</sup>, Zia-Ur-Rehman, M.<sup>c</sup>, Shafiq, M.<sup>b</sup>, Asiri, A.M.<sup>d</sup>

**Methyl 2-benzyl-4-hydroxy-1,1-dioxo-1,2,3,4-tetrahydro- $\lambda$ 6,2-benzothiazine-3-carboxyl-ate**

(2011) *Acta Crystallographica Section E: Structure Reports Online*, 67 (7), pp. o1588-o1589.

<sup>a</sup> Department of Physics, School of Physical Sciences, University of the Punjab, Lahore 54590, Pakistan

<sup>b</sup> Materials Chemistry Laboratory, Department of Chemistry, GC University, Lahore 54000, Pakistan

<sup>c</sup> Applied Chemistry Research Centre, PCSIR Laboratories Complex, Lahore 54600, Pakistan

<sup>d</sup> Center of Excellence for Advanced Materials Research, King Abdul Aziz University, PO Box 80203, Jeddah, Saudi Arabia

#### Abstract

In the title compound, C<sub>17</sub>H<sub>15</sub>NO<sub>5</sub>S, the benzene ring of the fused-ring system is twisted by 11.67 (6)° with respect to the thiazine ring. The atoms of the four-atom methyl ester group and the phenyl ring of the benzyl unit are inclined at 16.50 (7) and 44.52 (3)° with respect to the thiazine ring. An intra-molecular O - H...O hydrogen bond gives rise to a six-membered S(6) ring motif. In the crystal, molecules are extended through a C - H...O inter-action along the a axis. C - H... $\pi$  inter-actions are also observed. © Arshad et al. 2011.

**Document Type:** Article