

KINGDOM OF SAUDI ARABIA
Ministry of Higher Education
KING ABDULAZIZ UNIVERSITY
SCIENTIFIC COUNCIL
Scientific Promotion Committee



المملكة العربية السعودية
وزارة التعليم العالي
جامعة الملك عبد العزيز
المجلس العلمي
لجنة الترقية العلمية

Ref.

Date

الرقم

التاريخ ١٤٢٢/٩/١٣

إفادة

لتحديد الباحث الرئيسي في بحث مشترك مقدم للترقية العلمية

عنوان البحث : A FACILE SYNTHESIS of 8-SUBSTITUTED-4-

HYDROXY-2-METHYL-6,7-METHYLENEDI OXY-1,2,3,4-TETRAHYDROISOUQUINOLINES.

المؤلفون حسب ترتيب ظهورهم على البحث : د/ طارق رشاد سبكي ،
د. سالم أحمد ياسين ، د. عبد الله محمد عيسى ، د. محمد علي حسنة

مشور

مقبول للنشر ✓

جهة وتاريخ النشر مجلة المحرقة الكيميائية السعودية JSCS ١٤٢٢/٩/١٠
١٤٢٢/١١/٢٥

نشهد نحن المؤلفون المشاركون في البحث المذكور أعلاه - بأن الباحث الرئيسي لهذا البحث هو :

د/ طارق رشاد سبكي وهذه إفادة منا بذلك لتقديمها إلى لجنة الترقية العلمية .

د. عبد الله عيسى

أعضاء فريق البحث :

د. محمد علي حسنة

د. سالم أحمد ياسين

Attestation

The co-authors of the paper entitled A.FACILE SYNTHESIS of 8-SUBSTITUTED-4-

HYDROXY-2-METHYL-6,7-METHYLENEDI OXY-1,2,3,4-TETRAHYDROISOUQUINOLINES.

attest that Dr. Tariq R. Sobahi.

is the principal Investigator of this paper.

(Signature)

M. A. Hassan

A.M. ASIRI

نموذج رقم (١)

A FACILE SYNTHESIS OF 8-SUBSTITUTED-4-HYDROXY-2-METHYL-6,7-METHYLENEDIOXY-1,2,3,4-TETRAHYDROISOQUINOLINES

Tariq R. Sobahi, Salem A. Basaif, Abdullah M. Asiri and Mohamed A. Hassan

Department of Chemistry, Faculty of Science, King Abdulaziz University,
Jeddah-21589, P.O.Box 80203, Saudi Arabia

(Received 29th April 2001; Accepted 16th July 2001)

بسبب وجود التأثير المتلازم والمتناسق لمجموعتي البنزاييل أمينو و الألكوكسي المتجاورتين في مركب *N*-methyl-*N*-(2,2-diethoxyethyl)piperonylamine (3) ، الذي تم استخدامه كمادة أولية لتحضير عديد من مشتقات أيزوكينولين المستبدلة في الوضع -8 التي من المتوقع أن يكون لها نشاط بيولوجي فعال. يتم إحلال الليثيوم بسرعة و بانتقائية مرتفعة في المركب 3 عند الموقع ٢- بواسطة *n*-butyllithium في ثنائي إيثيل إيثير عند درجة صفر مئوية والذي يتفاعل بدوره مع الكتروفيلات المختلفة ومن ثم يتم الحولقة في وسط حمضي ليعطي سلسلة من المركبات 8-substituted-4-hydroxy-2-methyl-6,7-methylenedioxy-1,2,3,4-tetrahydroisoquinoline

Owing to the concomitant coordinating effect of the adjacent benzylamino and alkoxy groups, *N*-methyl-*N*-(2,2-diethoxyethyl)piperonylamine (3), which was used as a starting material for the synthesis of different 8-substituted-4-hydroxy-2-methyl-6,7-methylenedioxy-1,2,3,4-tetrahydroisoquinolines, is rapidly and exclusively lithiated at the position-2 with *n*-butyllithium in diethyl ether at 0°C. Successive reactions with electrophiles followed with cyclization in acidic medium give the corresponding 8-substituted-4-hydroxy-2-methyl-6,7-methylenedioxy-1,2,3,4-tetrahydroisoquinolines.

INTRODUCTION

The isoquinoline alkaloids are an important class of natural products including simple isoquinolines, like anhalamine produced by the cactus peyote, or benzoisoquinolines, such as morphine and codeine, present in the opiate poppies, whose powerful physiologic effects such as an anesthetics and analgesic effects are well-known. Probably due to this "bewitching" characteristics many chemists have attempted to synthesis these natural products as well as the synthesis of their analogues with the aim to modify their physiological activity [1-3].

It is well-known that different 8-substituted isoquinolines are rare due to their synthetic difficulties. In this general context we are engaged in the synthesis of 8-substituted-isoquinolines via lithiation of *N*-methyl-*N*-(2,2-diethoxyethyl)-piperonyl amine followed by ring closure in acid medium.

EXPERIMENTAL

Reagents: piperonal, bromoacetaldehyde diethyl acetal, dibenzyl disulfide, oxirane, paraformaldehyde and *n*-butyllithium (1.68M in hexane) were supplied by the Aldrich Chemical company, whilst *p*-toluenesulfonylazide was prepared by treatment of *p*-toluenesulfonylchloride with sodium azide.

Preparation of *N*-methylpiperonylamine (2).

Piperonal (1) (15g, 0.1mol) was added to 40% aqueous solution of methylamine, and the mixture stirred at room temperature for 1 hour. The crude reaction product was then extracted with ether (3x30ml). The combined ether extracts were washed with water (2x20ml), dried over MgSO₄, filtered and evaporated to yield the residue which was added to a solution of NaBH₄ (6.4g, 0.2mol) in anhydrous diethyl ether (50ml) while stirring under reflux for 1 hour. After cooling at 0°C, water (50ml) was cautiously added, the mixture filtered and the solid product was dissolved in chloroform (100ml) then washed

4.67 (1H, t, $J=5.6$ Hz, $CHCH_2$), 4.65 (2H, broad s, NH_2), 3.64 (2H, dq, $J=9.5, 7.1$ Hz, CH_2), 3.50 (2H, dq, $J=9.5, 7.1$ Hz, CH_2), superimposed on 3.50 (2H, s, CH_2N), 2.55 (2H, d, $J=5.6$ Hz, CH_2CH), 2.20 (3H, s, CH_3), 1.20 (6H, t, $J=7.0$ Hz, $2CH_3$) ppm; $C_{15}H_{24}N_2O_4$ (296), Calcd.: C, 60.81; H, 8.11; N, 9.46. Found: C, 60.67; H, 8.13; N, 9.35.

General procedure for cyclization of 4-6: Formation of 8-substituted-4-hydroxy-2-methyl-6,7-methylenedioxy-1,2,3,4-tetrahydroisoquinolines 8-10.

The crude products of 4-6 were treated, individually, with 20% aqueous hydrochloric acid (150ml) and allowed to react at room temperature for 16 hrs with stirring. The resulting solution was treated with charcoal, filtered and basified with 40% aqueous sodium hydroxide, while keeping the temperature below 40°C. The aqueous solution was extracted with dichloromethane (3x100ml). The collected organic phase was dried over anhydrous sodium sulfate, filtered and the solvent was evaporated *in vacuo* to give the desired 8-substituted isoquinolines 8-10.

Compound 8: Pale yellow crystals from ethyl acetate, mp 194-195°C; 1H NMR (200MHz, $CDCl_3$) δ 7.10 (5H, m, 5CH Ar), 6.79 (1H, s, CH Ar), 5.96 (1H, d, $J=1.3$ Hz, OCH_2O), 5.94 (1H, d, $J=1.3$ Hz, OCH_2O), 4.37 (1H, broad s, CHOH), 3.94 (2H, s, S- CH_2), 3.88 (1H, broad s, OH), 3.58 (1H, d, $J=15.7$ Hz, CH_2N), 2.82 (1H, ddd, $J=11.7, 1.7, 0.9$ Hz, CH_2CH), 2.64 (1H, d, $J=15.7$ Hz, CH_2N), 2.31 (3H, s, CH_3), 2.26 (1H, dd, $J=11.7, 1.7$ Hz, CH_2CH) ppm; $C_{18}H_{19}NO_5S$ (329), Calcd.: C, 65.63; H, 5.81; N, 4.25. Found: C, 65.47; H, 5.73; N, 4.37.

Compound 9: colorless crystals from ethanol, mp 182-183°C; 1H NMR (200MHz, $DMSO-d_6$) δ 6.80 (1H, s, CH Ar), 5.92 (1H, s, OCH_2O), 5.91 (1H, s, OCH_2O), 5.10 (1H, d, $J=7.0$ Hz, CHOH), 4.73 (1H, t, $J=5.4$ Hz, CH_2OH), 4.51 (1H, sym. m, $CHCH_2$), 3.50-3.30 (3H, m, $CH_2OH+CH_2CH_2N$), 3.29 (1H, d, $J=14.7$ Hz, CH_2CH_2N), 2.75 (1H, dd, $J=10.9, 5.0$ Hz, CH_2CH), 2.64 (2H, t, $J=7.4$ Hz, CH_2CH_2OH), 2.35 (3H, s, CH_3), 2.29 (1H, dd, $J=10.9, 7.3$ Hz, CH_2CH) ppm; $C_{13}H_{17}NO_4$ (251), Calcd.: C, 62.14; H, 6.82; N, 5.57. Found: C, 62.31; H, 6.74; N, 5.79.

Compound 10: colorless crystals from ligroin: ethanol (1:1), mp 174-175°C; 1H NMR (200MHz, $DMSO-d_6$) δ 6.86 (1H, s, CH Ar), 5.95 (1H, s,

OCH_2O), 5.94 (1H, s, OCH_2O), 5.14 (1H, d, $J=6.5$ Hz, CHOH), 4.94 (1H, t, $J=5.0$ Hz, CH_2OH), 4.52 (1H, sym. m, $CHCH_2$), 4.39 (2H, d, $J=5.0$ Hz, CH_2OH), 3.62 (1H, d, $J=15.1$ Hz, CH_2N), 3.36 (1H, d, $J=15.1$ Hz, CH_2N), 2.78 (1H, dd, $J=10.8, 5.3$ Hz, CH_2CH), 2.34 (3H, s, CH_3), 2.28 (1H, dd, $J=10.8, 7.3$ Hz, CH_2CH) ppm; $C_{12}H_{15}NO_4$ (237), Calcd.: C, 60.56; H, 6.37; N, 5.90. Found: C, 60.71; H, 6.19; N, 6.02.

RESULTS AND DISCUSSION


In order to synthesise different 8-substituted-4-hydroxy-2-methyl-6,7-methylenedioxy-1,2,3,4-tetrahydroisoquinolines, *N*-methyl-*N*-(2,2-diethoxyethyl)piperonylamine (3) was used as a starting material. Preparation of *N*-methyl-*N*-(2,2-diethoxyethyl)piperonylamine (3) was achieved by treatment of *N*-methyl-piperonylamine (2), which was prepared by reacting 40% aqueous methylamine with piperonal (1) and successively adding of sodium borohydride, bromoacetaldehyde diethyl acetal and aqueous potassium hydroxide solution at 100°C to produce the desired product 3 in very good yield 88% (Scheme 1).

Owing to the concomitant coordinating effect of the adjacent benzylamino and alkoxy groups, *N*-methyl-*N*-(2,2-diethoxyethyl)piperonylamine (3) is rapidly and exclusively lithiated at the position-2 with *n*-butyllithium in diethyl ether at 0°C [4-6]. Successive reactions with different electrophiles such as dibenzyl disulfide, oxirane, paraformaldehyde and *p*-toluenesulfonylazide give the corresponding substituted products 4-7, respectively (Scheme 2) [7,8].

Acid-catalyzed cyclization of the products 4-6 give access to a series of 8-substituted-4-hydroxy-2-methyl-6,7-methylenedioxy-1,2,3,4-tetrahydroisoquinolines 8-10, respectively, as shown in scheme 3 [9,10].

Reduction of the product *N*-methyl-*N*-(2,2-diethoxyethyl)-2-azido-3,4-methylenedioxybenzylamine (7) with lithium aluminium hydride in anhydrous diethyl ether under reflux, under nitrogen, produced *N*-methyl-*N*-(2,2-diethoxyethyl)-2-amino-3,4-methylenedioxybenzylamine (11) in excellent yield 95% (Scheme 4).

Unfortunately, cyclization of each azido- or amino-substituted products 7 and 11 respectively, gave a mixture of unidentified products.



Scheme 4

REFERENCES

- | | |
|--|---|
| <p>[1] B. Gøber, Arch Pharm. Weinheim, 299, 196 (1966).</p> <p>[2] P. Beak, Accounts Chem. Res., 19, 356 (1986).</p> <p>[3] P. Beak, Accounts Chem. Res., 15, 306 (1982).</p> <p>[4] N.S. Narasimhan, Synthesis, 957 (1983).</p> <p>[5] G. Simig, Synlett, 50 (1990).</p> | <p>[6] G. Simig and M. Schlosser, Tetrahedron Lett., 31, 3125 (1990).</p> <p>[7] V. Snieckus, Heterocycles, 14, 1649 (1980).</p> <p>[8] J.M. Bobbitt, Adv. Heterocyclic Chem., 15, 99 (1973).</p> <p>[9] D.L. Comins, J. Org. Chem., 54, 3730 (1989).</p> <p>[10] H.W. Gschwend, Org. Reactions, 26, 1 (1979).</p> |
|--|---|